

# The Cesàro $\chi^2$ of Tensor products in Orlicz sequence spaces

Research Article

Vandana<sup>a</sup>, Deepmala<sup>b</sup>, N. Subramanian<sup>c</sup>, Lakshmi Narayan Mishra<sup>d</sup>, \*<sup>a</sup> Department of Management Studies, Indian Institute of Technology, Madras, Chennai -600 036, Tamil Nadu, India<sup>b</sup> Mathematics Discipline, PDPM-Indian Institute of Information Technology, Design and Manufacturing, Jabalpur - 482005 (MP), India<sup>c</sup> Department of Mathematics, SASTRA University, Thanjavur-613 401, India<sup>d</sup> Department of Mathematics, School of Advanced Sciences, Vellore Institute of Technology (VIT) University, Vellore 632 014, Tamil Nadu, India

Received 02 March 2018; accepted (in revised version) 19 July 2018

**Abstract:** Let  $X$  be a Banach lattice and  $\chi_f^2$  be an double gai Orlicz sequence space associated to an Orlicz function with the  $\Delta_2$ -condition. In this paper we define the Cesàro  $\chi^2$  sequence space  $\text{Ces}_p^q(\chi_f^2)$  generated by a Orlicz sequence space and exhibit some general properties of the spaces.

**MSC:** 46B42 • 46B28

**Keywords:** Analytic sequence • Double sequences •  $\chi^2$  space • Cesàro  $\chi^2$  • Musielak-Orlicz function •  $p$ - metric space • Banach metric lattice • Positive tensor product

© 2018 The Author(s). This is an open access article under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/3.0/>).

## 1. Introduction

Throughout  $w, \chi$  and  $\Lambda$  denote the classes of all, gai and analytic scalar valued single sequences, respectively. We write  $w^2$  for the set of all complex double sequences  $(x_{mn})$ , where  $m, n \in \mathbb{N}$ , the set of positive integers. Then,  $w^2$  is a linear space under the coordinate wise addition and scalar multiplication.

Some initial works on double sequence spaces is found in Bromwich [1]. Later on it was investigated by Hardy [2], Moricz [3], Moricz and Rhoades [4], Basarir and Solankan [5], Tripathy et al. ([6] -[17]), Turkmenoglu [18], Raj ([19]-[25]), Mishra et al. ([29]-[33]) and many others. Let  $(x_{mn})$  be a double sequence of real or complex numbers. Then the series  $\sum_{m,n=1}^{\infty} x_{mn}$  is called a double series. The double series  $\sum_{m,n=1}^{\infty} x_{mn}$  give one space is said to be convergent if and only if the double sequence  $(S_{mn})$  is convergent, where

$$S_{mn} = \sum_{i,j=1}^{m,n} x_{ij} (m, n = 1, 2, 3, \dots).$$

A double sequence  $x = (x_{mn})$  is said to be double analytic if

$$\sup_{m,n} |x_{mn}|^{\frac{1}{m+n}} < \infty.$$

The vector space of all double analytic sequences are usually denoted by  $\Lambda^2$ . A sequence  $x = (x_{mn})$  is called double entire sequence if

\* Corresponding author.

E-mail address(es): [vdrai1988@gmail.com](mailto:vdrai1988@gmail.com) (Vandana), [dmrai23@gmail.com](mailto:dmrai23@gmail.com) (Deepmala), [nsmaths@yahoo.com](mailto:nsmaths@yahoo.com) (N. Subramanian), [lakshminarayanmishra04@gmail.com](mailto:lakshminarayanmishra04@gmail.com) (Lakshmi Narayan Mishra).

$$|x_{mn}|^{\frac{1}{m+n}} \rightarrow 0 \text{ as } m, n \rightarrow \infty.$$

The vector space of all double entire sequences are usually denoted by  $\Gamma^2$ . Let the set of sequences with this property be denoted by  $\Lambda^2$  and  $\Gamma^2$  is a metric space with the metric

$$d(x, y) = \sup_{m, n} \left\{ |x_{mn} - y_{mn}|^{\frac{1}{m+n}} : m, n : 1, 2, 3, \dots \right\}, \quad (1)$$

for all  $x = \{x_{mn}\}$  and  $y = \{y_{mn}\}$  in  $\Gamma^2$ . Let  $\phi = \{\text{finite sequences}\}$ .

Consider a double sequence  $x = (x_{mn})$ . The  $(m, n)^{th}$  section  $x^{[m, n]}$  of the sequence is defined by  $x^{[m, n]} = \sum_{i, j=0}^{m, n} x_{ij} \delta_{ij}$  for all  $m, n \in \mathbb{N}$ ,

$$\delta_{mn} = \begin{pmatrix} 0 & 0 & \dots & 0 & 0 & \dots \\ 0 & 0 & \dots & 0 & 0 & \dots \\ \cdot & & & & & \\ \cdot & & & & & \\ \cdot & & & & & \\ 0 & 0 & \dots & 1 & 0 & \dots \\ 0 & 0 & \dots & 0 & 0 & \dots \end{pmatrix}$$

with 1 in the  $(m, n)^{th}$  position and zero otherwise.

A double sequence  $x = (x_{mn})$  is called double gai sequence if  $((m+n)! |x_{mn}|)^{\frac{1}{m+n}} \rightarrow 0$  as  $m, n \rightarrow \infty$ . The double gai sequences will be denoted by  $\chi^2$ .

## 2. Definitions and Preliminaries

### Definition 2.1 (26).

An Orlicz function is a function  $M : [0, \infty) \rightarrow [0, \infty)$  which is continuous, non-decreasing and convex with  $M(0) = 0$ ,  $M(x) > 0$ , for  $x > 0$  and  $M(x) \rightarrow \infty$  as  $x \rightarrow \infty$ . If convexity of Orlicz function  $M$  is replaced by  $M(x+y) \leq M(x) + M(y)$ , then this function is called modulus function. An Orlicz function  $M$  is said to satisfy  $\Delta_2$ -condition for all values  $u$ , if there exists  $K > 0$  such that  $M(2u) \leq KM(u)$ ,  $u \geq 0$ .

### Lemma 2.1.

Let  $M$  be an Orlicz function which satisfies  $\Delta_2$ -condition and let  $0 < \delta < 1$ . Then for each  $t \geq \delta$ , we have  $M(t) < K\delta^{-1}M(2)$  for some constant  $K > 0$ .

### Definition 2.2 (27).

Let  $n \in \mathbb{N}$  and  $X$  be a real vector space of dimension  $m$ , where  $n \leq m$ . A real valued function  $d_p(x_1, \dots, x_n) = \|(d_1(x_1, 0), \dots, d_n(x_n, 0))\|_p$  on  $X$  satisfying the following four conditions:

(i)  $\|(d_1(x_1, 0), \dots, d_n(x_n, 0))\|_p = 0$  if and only if  $d_1(x_1, 0), \dots, d_n(x_n, 0)$  are linearly dependent,

(ii)  $\|(d_1(x_1, 0), \dots, d_n(x_n, 0))\|_p$  is invariant under permutation,

(iii)  $\|(\alpha d_1(x_1, 0), \dots, \alpha d_n(x_n, 0))\|_p = |\alpha| \|(d_1(x_1, 0), \dots, d_n(x_n, 0))\|_p$ ,  $\alpha \in \mathbb{R}$

(iv)  $d_p((x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)) = (d_X(x_1, x_2, \dots, x_n)^p + d_Y(y_1, y_2, \dots, y_n)^p)^{1/p}$  for  $1 \leq p < \infty$ ; (or)

(v)  $d((x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)) := \sup \{d_X(x_1, x_2, \dots, x_n), d_Y(y_1, y_2, \dots, y_n)\}$ ,

for  $x_1, x_2, \dots, x_n \in X$ ,  $y_1, y_2, \dots, y_n \in Y$  is called the  $p$  product metric of the Cartesian product of  $n$  metric spaces is the  $p$  norm of the  $n$ -vector of the norms of the  $n$  subspaces.

A trivial example of  $p$  product metric of  $n$  metric space is the  $p$  norm space is  $X = \mathbb{R}$  equipped with the following Euclidean metric in the product space is the  $p$  norm:

$$\|(d_1(x_1, 0), \dots, d_n(x_n, 0))\|_E = \sup(|\det(d_{mn}(x_{mn}, 0))|) = \sup \left( \begin{pmatrix} d_{11}(x_{11}, 0) & d_{12}(x_{12}, 0) & \dots & d_{1n}(x_{1n}, 0) \\ d_{21}(x_{21}, 0) & d_{22}(x_{22}, 0) & \dots & d_{2n}(x_{2n}, 0) \\ \cdot & & & \\ \cdot & & & \\ \cdot & & & \\ d_{n1}(x_{n1}, 0) & d_{n2}(x_{n2}, 0) & \dots & d_{nn}(x_{nn}, 0) \end{pmatrix} \right)$$

where  $x_i = (x_{i1}, \dots, x_{in}) \in \mathbb{R}^n$  for each  $i = 1, 2, \dots, n$ .

If every Cauchy sequence in  $X$  converges to some  $L \in X$ , then  $X$  is said to be complete with respect to the  $p$ - metric. Any complete  $p$ - metric space is said to be  $p$ - Banach metric space.

Let  $M$  and  $\Phi$  be mutually complementary Orlicz functions. Then, we have

(i) For all  $u, y \geq 0$ ,

$$uy \leq M(u) + \Phi(y), \text{ (Young's inequality) [See [Kampthorn et al. , [28]]} \tag{2}$$

(ii) For all  $u \geq 0$ ,

$$u\eta(u) = M(u) + \Phi(\eta(u)). \tag{3}$$

(iii) For all  $u \geq 0$ , and  $0 < \lambda < 1$ ,

$$M(\lambda u) \leq \lambda M(u). \tag{4}$$

A sequence  $M = (M_{mn})$  of Orlicz function is called a Musielak-Orlicz function. A sequence  $g = (g_{mn})$  defined by

$$g_{mn}(v) = \sup \{ |v| u - M_{mn}(u) : u \geq 0 \}, m, n = 1, 2, \dots$$

is called the complementary function of a Musielak-Orlicz function  $M$ . For a given Musielak Orlicz function  $M$ , the Musielak-Orlicz sequence space  $t_f$  is defined by

$$t_f = \{ x \in w^2 : I_f(|x_{mn}|)^{1/m+n} \rightarrow 0 \text{ as } m, n \rightarrow \infty \}.$$

We consider  $t_f$  equipped with the Luxemburg metric

$$d(x, y) = \sup_{mn} \left\{ \inf \left( \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} f_{mn} \left( \frac{|x_{mn}|^{1/m+n}}{mn} \right) \right) \leq 1 \right\}.$$

The positivity perspective, it is known that the projective tensor and the injective tensor product of two Banach lattices are, in general not Banach lattices. Throughout  $w, \chi$  and  $\Lambda$  denote the classes of all, gai and analytic scalar valued single sequences, respectively. We write  $w^2$  for the set of all complex double sequences  $(x_{mn})$ , where  $m, n \in \mathbb{N}$ , the set of positive integers. Then,  $w^2$  is a linear space under the coordinate wise addition and scalar multiplication.

**Definition 2.3.**

**Positive tensor products:** For Banach lattices  $X$  and  $Y$ , let  $X \otimes Y$  denote the algebraic tensor product of  $X$  and  $Y$ . For each

$$u = \sum_{m=1}^r \sum_{n=1}^s x_{mn} \otimes y_{mn} \in X \otimes Y,$$

define  $T_u : X^* \rightarrow Y$  by  $T_u(x^*) = \sum_{m=1}^r \sum_{n=1}^s x^*(x_{mn}) y_{mn}$  for each  $x^* \in X^*$ . Then injective cone on  $X \otimes Y$  is defined to be  $C_i = \{ u \in X \otimes Y : T_u(x^*) \in Y^* \ \forall x^* \in X^{**} \}$ .

**Definition 2.4.**

Let  $X \bar{\otimes}_i Y$  denote the completion of  $X \otimes Y$  with respect  $d(.,.)$ . Then  $X \bar{\otimes}_i Y$  with  $C_i$  as its positive cone is a Banach lattice called the positive injective tensor product of  $X$  and  $Y$ .

The positive cone on  $x \otimes Y$  is defined to be

$$C_p = \{ \sum_{m=1}^r \sum_{n=1}^s x_{mn} \otimes y_{mn} : r, s \in \mathbb{N}, x_{mn} \in X^+, y_{mn} \in Y^+ \}.$$

We define the following spaces:

For a Banach metric lattice  $X$ , let

$$\chi_f^2(X) = \left\{ \bar{x} = (x_{ij})_{ij} \in X^{\mathbb{N} \times \mathbb{N}} : \left( x^* \left( (i+j)! |x_{ij}| \right)^{1/i+j} \right)_{ij} \in \chi^2, \forall x^* \in X^{**} \right\}.$$

The metric defined to be

$$d(x, y) = \sup \left\{ \left\| \left( x^* \left( (i+j)! |x_{ij} - y_{ij}| \right)^{1/i+j} \right) : x^* \in B_{X^{**}} \right\| \right\}, x = (x_{ij})_{ij} \in \chi_f^2(X).$$

Let  $\chi_f^3(X) = \{ \bar{x} \in \chi_f^2(X) : \lim_{i,j,n} \left\| \left( (i+j)! |\bar{x}_{ij}(>n)| \right)^{1/i+j} \right\| \rightarrow 0 \text{ as } i, j \rightarrow \infty \}$ , with the metric

$$d(x, y) = \sup \left\{ \left\| \left( (i+j)! |(\bar{x}_{ij} - \bar{y}_{ij})(>n)| \right)^{1/i+j} \right\| \forall (\bar{x}_{ij}) \in \chi_f^2(X^{**}) \right\}.$$

### 3. Notations

For a vector space  $X$ , a vector  $\bar{x} = (x_{ij})_{ij} \in X^{\mathbb{N} \times \mathbb{N}}$  and  $n \in \mathbb{N}$ , we write  $\bar{x}(\leq n)$  is a two dimensional matrix from first term to  $n$ th term and remaining term zero. and  $\bar{x}(> n)$  is a two dimensional matrix from first term to  $n$ th term zero and start with  $(n+1)$ th term.

If  $X$  is an ordered set, the usual order on  $X^{\mathbb{N} \times \mathbb{N}}$  is defined by  $\bar{x} = (x_{ij})_{ij} \geq 0 \iff (x_{ij}) \geq 0$  for each  $i, j \in \mathbb{N}$ . for Banach lattice  $X$ ,  $X^*$  denotes its dual space,  $B_X$  denotes its closed unit ball, and  $X^+$  denotes its positive cone.

### 4. Some New Cesàro Orlicz sequence space of Tensor products in Musielak

The main aim of this article is to introduce the following sequence spaces and examine the topological and algebraic properties of the resulting sequence spaces. Let  $f = (f_{mn})$  be a sequence Orlicz function,  $(\bar{X}, \|(d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p)$  be a  $p$ - metric space, and consider  $\mu_{mn}(\bar{x}) = \left\| \left| ((i+j)! |\bar{x}_{ij}(>n)|)^{1/i+j} \right| \right\|$  and  $\eta_{mn}(\bar{x}) = \left\| \left| |\bar{x}_{ij}(>n)|^{1/i+j} \right| \right\|$ .

We define the following sequence spaces as follows:

$$\left[ \chi_f^2, \|(d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right] = \lim_{m,n \rightarrow \infty} \left\{ \sum_m \sum_n \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right] = 0 \right\},$$

and

$$\left[ \Lambda_f^2, \|(d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right] = \sup \left\{ \sum_m \sum_n \left[ f_{mn} \left( \|\eta_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right] < \infty \right\}.$$

Let  $p \in [1, \infty)$  and  $q$  be a double gai sequence of positive real numbers such that

$$Q_{ij} = \sum_{m=0}^i \sum_{n=0}^j q_{mn}, i, j \in \mathbb{N}$$

$$Ces_p^q(\chi_f^2) = d(x, 0) =$$

$$x \in \chi^2 := \lim_{m,n \rightarrow \infty}$$

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{p_{mn}} \right]^{1/p_{mn}}$$

$= 0$

If  $q_{mn} = 1$  for all  $m, n \in \mathbb{N}$ , then  $Ces_p^q(\chi_f^2)$  reduces to  $Ces_p(\chi_f^2)$  and if  $f(x) = x$  then  $Ces_p^q(\chi_f^2)$  reduces to  $Ces_p^q(\chi^2)$ .

Let  $p \in [1, \infty)$  and  $q$  be a double analytic sequence of positive real numbers such that

$$Q_{ij} = \sum_{m=0}^i \sum_{n=0}^j q_{mn}, i, j \in \mathbb{N}$$

$$Ces_p^q(\Lambda_f^2) = d(x, 0) =$$

$$x \in \Lambda^2 := \sup$$

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\eta_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{p_{mn}} \right)^{1/p_{mn}}$$

$< \infty$

If  $q_{mn} = 1$  for all  $m, n \in \mathbb{N}$ , then  $Ces_p^q(\Lambda_f^2)$  reduces to  $Ces_p(\Lambda_f^2)$  and if  $f(x) = x$  then  $Ces_p^q(\Lambda_f^2)$  reduces to  $Ces_p^q(\Lambda^2)$ .

The space  $Ces_p^q(\chi_f^2)$  is a metric space with the metric

$$d(x, y) = \inf$$

$$\sup \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn}$$

$$\left[ f_{mn} \|\mu_{mn}(\bar{x}) - 0, (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right]^{p_{mn}} \right]^{1/p_{mn}} \leq 1$$

The space  $Ces_p^q(\Lambda_f^2)$  is a metric space with the metric

$$d(x, y) = \inf$$

$$\sup \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn}$$

$$\left[ f_{mn} \left( \|\eta_{mn}(\bar{x}) - 0, (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{p_{mn}} \right]^{1/p_{mn}} \leq 1.$$

### 5. Main Results

#### Proposition 5.1.

Let, the tensor produce of Orlicz sequence space  $x, y \in Ces_p^q(\chi_f^2)$ . Then for any  $\epsilon > 0$  and  $L > 0$ , there exists  $\delta > 0$  such that  $(d(x + y, 0), 0)^{p_{mn}} = d(x, 0)^{p_{mn}} + \epsilon$ , whenever  $d(x, 0)^{p_{mn}} \leq L$  and  $d(y, 0)^{p_{mn}} \leq \delta$

*Proof.* For any fix  $\epsilon > 0$

$$\begin{aligned}
 & d(x + y, 0)^{p_{mn}} = \\
 & \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x} + \bar{y}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right] \right)^{p_{mn}} \\
 & \leq \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right] \right)^{p_{mn}} + \\
 & \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{y}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right] \right)^{p_{mn}} \leq (1 - \beta) \\
 & \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right] \right)^{p_{mn}} + \\
 & \beta \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right] \right)^{p_{mn}} \quad + \\
 & \beta \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \left\| \frac{\mu_{mn}(\bar{y})}{\beta}, (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0)) \right\|_p \right) \right] \right)^{p_{mn}} \\
 & \leq (1 - \beta) \\
 & \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right] \right)^{p_{mn}} \quad + \\
 & \frac{\beta}{2} \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j 2q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right] \right)^{p_{mn}} \\
 & + \frac{\beta}{2} \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j 2q_{mn} \left[ f_{mn} \left( \left\| \frac{\mu_{mn}(\bar{y})}{\beta}, (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0)) \right\|_p \right) \right] \right)^{p_{mn}} \\
 & \leq d(x, 0)^{p_{mn}} + \frac{\epsilon}{2} + \\
 & \left( \frac{2}{\beta} \right)^{p_{mn}-1} \\
 & \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right] \right)^{p_{mn}} \\
 & \leq d(x, 0)^{p_{mn}} + \frac{\epsilon}{2} + \frac{\epsilon}{2} \\
 & \leq d(x, 0)^{p_{mn}} + \epsilon. \quad \square
 \end{aligned}$$

#### Proposition 5.2.

For every  $p = (p_{mn})$ ,  $[Ces_p^q(\Lambda_f^2)]^\beta = [Ces_p^q(\Lambda_f^2)]^\alpha = [Ces_p^q(\Lambda_f^2)]^\gamma = [Ces_p^q(\eta_f^2)]^\beta$ , where  $[Ces_p^q(\eta_f^2)] = \bigcap_{N \in \mathbb{N} - \{1\}} \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}) N^{m+n/p_{mn}}, (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right] \right)^{p_{mn}} < \infty$

*Proof.* (1): First we show that  $[Ces_p^q(\eta_f^2)] \subset [Ces_p^q(\Lambda_f^2)]^\beta$

Let  $x \in [Ces_p^q(\eta_f^2)]$  and  $y \in [Ces_p^q(\Lambda_f^2)]^\beta$ . Then we can find a positive integer  $N$  such that  $(\|\mu(\bar{y})\|) < \max$

$$1, \sup_{m, n \geq 1} \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn}$$

$$\left[ f_{mn} \left( \|\mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{p_{mn}} < N, \text{ for all } m, n.$$

Hence we may write

$$|\sum_{m, n} \mu_{mn}(\bar{x}\bar{y})| \leq \sum_{m, n} |\mu_{mn}(\bar{x}\bar{y})| \leq \sum_{m, n} \mu_{mn}(\bar{x}) N^{m+n}.$$

Since  $x \in Ces_p^q(\eta_f^2)$ . the series on the right side of the above inequality is convergent, whence  $x \in Ces_p^q(\Lambda_f^2)$ . Hence

$$[Ces_p^q(\eta_f^2)] \subset [Ces_p^q(\Lambda_f^2)]^\beta$$

Now we show that  $[Ces_p^q(\Lambda_f^2)]^\beta \subset [Ces_p^q(\eta_f^2)]$

For this, let  $x \in [Ces_p^q(\Lambda_f^2)]^\beta$  and suppose that  $x \notin [Ces_p^q(\eta_f^2)]$ . Then there exists a positive integer  $N > 1$  such that  $\mu_{mn}(\bar{x}) N^{m+n} = \infty$ .

If we define  $\mu_{mn}(\bar{y}) = N^{m+n} \text{Sgn} \mu_{mn}(\bar{x})$   $m, n = 1, 2, \dots$ , then  $\mu_{mn}(\bar{y}) \in [Ces_p^q(\Lambda_f^2)]$ .

But, since

$|\sum_{m,n} \mu_{mn}(\bar{x}\bar{y})| = \sum_{m,n} |\mu_{mn}(\bar{x}\bar{y})| = \sum_{m,n} |\mu_{mn}(\bar{x})| N^{m+n} = \infty$ , we get  $x \notin [Ces_p^q(\Lambda_f^2)]^\beta$ , which contradicts to the assumption  $x \in [Ces_p^q(\Lambda_f^2)]^\beta$ . Therefore  $x \in [Ces_p^q(\eta_f^2)]$ . Therefore  $[Ces_p^q(\Lambda_f^2)]^\beta = [Ces_p^q(\eta_f^2)]$ .

(ii) and (iii) can be shown in a similar way of (i). Therefore we omit it.  $\square$

**Proposition 5.3.**

Let  $p = (p_{mn})$  be a tensor products of Cesàro space of double analytic Orlicz sequence of strictly positive real numbers  $p_{mn}$ . Then

(i)  $Ces_p^q(\Lambda_f^2)$  is a paranormed space with

$$g(x) = \sup \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left\| \mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0)) \right\|_p^{1/M} \right] \quad (5)$$

if and only if  $h = \inf p_{mn} > 0$ , where  $M = \max(1, H)$  and  $H = \sup p_{mn}$ .

(ii)  $Ces_p^q(\Lambda_f^2)$  is a complete paranormed linear metric space if the condition  $p$  in (5.1) is satisfied.

**Proof. (i): Sufficiency:** Let  $h > 0$ . It is trivial that  $g(\theta) = 0$  and  $g(-x) = g(x)$ .

The inequality  $g(x+y) \leq g(x) + g(y)$  follows from the inequality (5.1), since  $p_{mn}/M \leq 1$  for all positive integers  $m, n$ . We also may write  $g(\lambda x) \leq \max(|\lambda|, |\lambda|^{h/M}) g(x)$ , since  $|\lambda|^{p_{mn}} \leq \max(|\lambda|^h, |\lambda|^M)$  for all positive integers  $m, n$  and for any  $\lambda \in \mathbb{C}$ , the set of complex numbers. Using this inequality, it can be proved that  $\lambda x \rightarrow \theta$ , when  $x$  is fixed and  $\lambda \rightarrow 0$ , or  $\lambda \rightarrow 0$  and  $x \rightarrow \theta$ , or  $\lambda$  is fixed and  $x \rightarrow \theta$ .

**Necessity:** Let  $Ces_p^q(\Lambda_f^2)$  be a paranormed space with the paranorm

$g(x) = \sup \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \left\| \mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0)) \right\|_p \right)^{1/M} \right]$  and suppose that  $h = 0$ . Since  $|\lambda|^{p_{mn}/M} \leq |\lambda|^{h/M} = 1$  for all positive integers  $m, n$  and  $\lambda \in \mathbb{C}$  such that  $0 < |\lambda| \leq 1$ , we have

$$\sup \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \left\| \mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0)) \right\|_p \right)^{1/M} \right] = 1.$$

Hence it follows that

$g(\lambda x) = \sup \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \left\| \mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0)) \right\|_p \right)^{1/M} \right] = 1$  for  $x = (\alpha) \in Ces_p^q(\Lambda_f^2)$  as  $\lambda \rightarrow 0$ . But this contradicts the assumption  $Ces_p^q(\Lambda_f^2)$  is a paranormed space with  $g(x)$ .

(ii) The proof is clear.  $\square$

**Proposition 5.4.**

$Ces_p^q(\Lambda_f^2)$  is a complete paranormed space with the natural paranorm if and only if  $Ces_p^q(\Lambda_f^2) = Ces^q(\Lambda_f^2)$ .

**Proposition 5.5.**

For every  $p = (p_{mn})$ , then  $Ces_p^q(\eta_f^2) \subset [Ces_p^q(\chi_f^2)]^\beta \not\subset Ces_p^q(\Lambda_f^2)$

**Proof. Case (i)** First we show that  $Ces_p^q(\eta_f^2) \subset [Ces_p^q(\chi_f^2)]^\beta$ . We know that  $[Ces_p^q(\chi_f^2)] \subset Ces_p^q(\Lambda_f^2)$ .

$[Ces_p^q(\Lambda_f^2)]^\beta \subset [Ces_p^q(\chi_f^2)]^\beta$ . But  $[Ces_p^q(\Lambda_f^2)]^\beta = Ces_p^q(\eta_f^2)$ . Therefore

$$Ces_p^q(\eta_f^2) \subset [Ces_p^q(\chi_f^2)]^\beta \quad (6)$$

**Case (2):** Now we show that  $[Ces_p^q(\chi_f^2)]^\beta \not\subset Ces_p^q(\Lambda_f^2)$ .

Let  $y = \mu_{mn}(\bar{y})$  be an arbitrary point  $[Ces_p^q(\chi_f^2)]^\beta$ . If  $y$  is not  $Ces_p^q(\Lambda_f^2)$ , then for each natural number  $d$ , we can find an index  $m_d n_d$  such that

$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{m_d n_d} \left[ f_{m_d n_d} \left( \left\| \mu_{m_d n_d}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0)) \right\|_p \right)^{1/M} \right] > d, (1, 2, 3, \dots)$ . Define  $x =$

$\{x_{mn}\}$  by  
 $\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{m_d n_d}$   
 $\left[ f_{mn} \left( \|\mu_{m_d n_d}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{1/M} > d$ , for  $(mn) = (m_d n_d)$  for some  $d \in \mathbb{N}$ ; and  
 $\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{m_d n_d}$   
 $\left[ f_{mn} \left( \|\mu_{m_d n_d}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{1/M} = 0$ , otherwise. Then  $x$  is  $Ces_p^q(\chi_f^2)$ , but for infinitely  $mn$ ,

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{1/M} > 1, \tag{7}$$

Consider the sequence  $z = \{z_{mn}\}$ , where  $\mu_{mn}(\bar{z}) = \mu_{mn}(\bar{x}) - s$  with

$$s = \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{1/M};$$

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{z}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{1/M} =$$

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{1/M}$$

The  $z$  is a point of  $Ces_p^q(\chi_f^2)$ . Also

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{z}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{1/M} = 0. \text{ Hence } z \text{ is in } Ces_p^q(\chi_f^2).$$

But, by the equation (7),

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}\bar{z}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{1/M} \text{ does not converge.}$$

$\mu_{mn}(\bar{x}\bar{y})$  diverges. Thus, the sequence  $\mu_{mn}(\bar{y})$  would not be  $\left[ Ces_p^q(\chi_f^2) \right]^\beta$ . This contradiction proves that

$$\left[ Ces_p^q(\chi_f^2) \right]^\beta \subset Ces_p^q(\Lambda_f^2) \tag{8}$$

If we now choose  $f = id$ , where  $id$  is the identity and  $\frac{1}{Q_{ij}}(q_{1n}\mu_{1n}(\bar{y})) = \frac{1}{Q_{ij}}(q_{1n}\mu_{1n}(\bar{x}))$  and  $\frac{1}{Q_{ij}}(q_{1n}\mu_{mn}(\bar{y})) = \frac{1}{Q_{ij}}(q_{1n}\mu_{mn}(\bar{x})) = 0$  ( $m, i > 1$ ) for all  $n, j$ , then obviously  $x \in Ces_p^q(\chi_f^2)$  and  $y \in Ces_p^q(\Lambda_f^2)$ , but

$$\sum \sum \mu_{mn}(\bar{x}\bar{y}) = \infty. \text{ Hence } \mu_{mn}(\bar{y}) \notin \left[ Ces_p^q(\chi_f^2) \right]^\beta \tag{9}$$

From (8) and (9), we are granted  $\left[ Ces_p^q(\chi_f^2) \right]^\beta \subsetneq Ces_p^q(\Lambda_f^2)$ . □

**Proposition 5.6.**

In tensor product of Orlicz sequence space of  $Ces_p^q(\chi_f^2)$  weak convergence does not imply strongly convergence.

*Proof.* Assume that weak convergence implies strong convergence  $Ces_p^q(\chi_f^2)$ . Then we would have  $\left[ Ces_p^q(\chi_f^2) \right]^{\beta\beta} = Ces_p^q(\chi_f^2)$  [see Wilansky ]. But  $\left[ Ces_p^q(\chi_f^2) \right]^{\beta\beta} \subsetneq \left[ Ces_p^q(\Lambda_f^2) \right]^\beta = Ces_p^q(\eta_f^2)$ . Thus  $\left[ Ces_p^q(\chi_f^2) \right]^{\beta\beta} \neq Ces_p^q(\chi_f^2)$ . Hence tensor product of Orlicz sequence space is weak convergence does not imply strong convergence in  $Ces_p^q(\chi_f^2)$ .

**Proposition 5.7.**

Let tensor product of Orlicz sequence space of  $f$  satisfies the  $\Delta_2$ - condition. Then  $Ces_p^q(\chi^2) \subset Ces_p^q(\chi_f^2)$ .

**Proof:** Let

$$x \in Ces_p^q(\chi^2) \tag{10}$$

Then  $\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}\bar{z}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{1/M}$   
 $\leq \epsilon$  for sufficiently large  $m, n$  and every  $\epsilon > 0$ .

$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}\bar{z}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{1/M} \leq f(\epsilon)$  (because the tensor product of Orlicz sequence space of  $f$  is non-decreasing)

$\Rightarrow \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}\bar{z}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{1/M} \leq Kf(\epsilon) < \epsilon$  (by the tensor product of Orlicz sequence space of  $\Delta_2$ -condition, for some  $K > 0$  and by defining  $f(\epsilon) < \frac{\epsilon}{K}$ )

$$\lim_{m, n \rightarrow \infty} \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \|\mu_{mn}(\bar{x}\bar{z}), (d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0))\|_p \right) \right]^{1/M} = 0 \quad (11)$$

Hence

$$x \in Ces_p^q(\chi_f^2). \quad (12)$$

From (10) and (12) we get  $Ces_p^q(\chi^2) \subset Ces_p^q(\chi_f^2)$ .  $\square$

**Proposition 5.8.**

$$\left[ Ces_p^q(\Lambda_f^2) \right]^\beta \subsetneq Ces_p^q(\chi_f^2)$$

*Proof.* Let  $\mu_{mn}(\bar{x}) \in \left[ Ces_p^q(\Lambda_f^2) \right]^\beta$

$$\sum \sum \mu_{mn}(\bar{x}\bar{y}) < \infty \forall \mu_{mn}(\bar{y}) \in \left[ Ces_p^q(\Lambda_f^2) \right]^\beta \quad (13)$$

Assume that  $\mu_{mn}(\bar{x}) \notin Ces_p^q(\chi_f^2)$ . Then there exist a sequence positive integers

$$\mu_{m_r n_r}(\bar{x}) > \frac{1}{((m_r + n_r! 2)^{(m_r + n_r)})}, (r = 1, 2, 3, \dots)$$

Take

$$\mu_{m_r n_r}(\bar{y}) = (2(m_r + n_r))^{m_r + n_r} \text{ for } r = 1, 2, 3, \dots$$

$$\mu_{m_r n_r}(\bar{y}) = 0 \text{ otherwise.}$$

Then  $\mu_{mn}(\bar{y}) \in \left[ Ces_p^q(\Lambda_f^2) \right]^\beta$ . But

$$\sum \sum \mu_{mn}(\bar{x}\bar{y}) > 1 + 1 + 1 + \dots$$

We know that the infinite series  $1 + 1 + 1 + \dots$  diverges. Hence  $\sum \sum \mu_{mn}(\bar{x}\bar{y})$  diverges. Which is contradicts. Hence  $\mu_{mn}(\bar{x}) \in Ces_p^q(\chi_f^2)$ . Therefore

$$\left[ Ces_p^q(\Lambda_f^2) \right]^\beta \subset Ces_p^q(\chi_f^2) \quad (14)$$

If we now choose  $p = (p_{mn})$  is a constant  $f = id$ , where  $id$  is the identity and  $\frac{1}{Q_{1j}}(q_{1n}\mu_{1n}(\bar{y})) = \frac{1}{Q_{1j}}(q_{1n}\mu_{1n}(\bar{x}))$  and  $\frac{1}{Q_{1j}}(q_{1n}\mu_{mn}(\bar{y})) = \frac{1}{Q_{1j}}(q_{1n}\mu_{mn}(\bar{x})) = 0$  ( $m, i > 1$ ) for all  $n, j$ , then obviously  $\mu_{mn}(\bar{x}) \in Ces_p^q(\chi_f^2)$  and  $\mu_{mn}(\bar{y}) \in Ces_p^q(\Lambda_f^2)$ , but

$$\sum \sum \mu_{mn}(\bar{x}\bar{y}) = \infty. \text{ Hence } \mu_{mn}(\bar{y}) \notin \left[ Ces_p^q(\chi_f^2) \right]^\beta \quad (15)$$

From (14) and (15) we are granted  $\left[ Ces_p^q(\Lambda_f^2) \right]^\beta \subsetneq Ces_p^q(\chi_f^2)$ .  $\square$

**Proposition 5.9.**

Let  $\left( Ces_p^q(\chi_f^2) \right)^*$  denote the dual space of  $Ces_p^q(\chi_f^2)$ . Then we have  $\left( Ces_p^q(\chi_f^2) \right)^* = Ces_p^q(\Lambda_f^2)$ .

*Proof.* We recall that



$$x = \mathfrak{S}_{mn} = \begin{pmatrix} 0 & 0 & \dots & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 & \dots & 0 \\ \vdots & & & & & \\ 0 & 0 & \dots & \frac{1}{(m+n)!} & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 & \dots & 0 \end{pmatrix}.$$

with  $\frac{1}{(m+n)!}$  in the  $(m, n)^{th}$  position and zero other wise, with

$$x = \mathfrak{S}_{mn}, \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} \left[ f_{mn} \left( \left\| \mu_{mn}(\bar{x}\bar{z}), d(\bar{x}_1, 0), d(\bar{x}_2, 0), \dots, d(\bar{x}_{n-1}, 0) \right\|_p \right) \right]^{1/M} \right) = \begin{pmatrix} 0 & 0 & \dots & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 & \dots & 0 \\ \vdots & & & & & \\ 0 & 0 & \dots & \frac{(m+n)!}{(m+n)!} & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 & \dots & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & \dots & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 & \dots & 0 \\ \vdots & & & & & \\ 0 & 0 & \dots & 1^{1/m+n} & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 & \dots & 0 \end{pmatrix}.$$

which is a  $Ces_p^q(\chi_f^2)$  sequence . Hence  $\mathfrak{S}_{mn} \in Ces_p^q(\chi_f^2)$ . Let us take  $f(x) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \mu_{mn}(\bar{x}\bar{y})$  with  $\mu_{mn}(\bar{x}) \in Ces_p^q(\chi_f^2)$ . and  $f \in (Ces_p^q(\chi_f^2))^*$ . Take  $x = \mu_{mn}(\bar{x}) = \mathfrak{S}_{mn} \in Ces_p^q(\chi_f^2)$ . Then

$$\mu_{mn}(\bar{y}) \leq \|f\| d(\mathfrak{S}_{mn}, 0) < \infty \text{ for each } m, n$$

Thus  $\mu_{mn}(\bar{y})$  is a bounded sequence and hence an Cesàro double analytic sequence of modulus. In other words  $\mu_{mn}(\bar{y}) \in Ces_p^q(\Lambda_f^2)$ . Therefore  $(Ces_p^q(\chi_f^2))^* = Ces_p^q(\Lambda_f^2)$ . □

### Competing Interests

The authors declare that there is no conflict of interests regarding the publication of this research paper.

### Acknowledgement

The authors are extremely grateful to the anonymous learned referee(s) for their keen reading, valuable suggestion and constructive comments for the improvement of the manuscript. The authors are thankful to the editor(s) and reviewers of International Journal of Advances in Applied Mathematics and Mechanics.

### References

- [1] T.J.I.A. Bromwich, An introduction to the theory of infinite series, Macmillan and Co.Ltd. New York, 1965.
- [2] G.H. Hardy, On the convergence of certain multiple series, Proceedings of the Cambridge Philosophical Society 19 (1917) 86-95.
- [3] F. Moricz, Extensions of the spaces  $c$  and  $c_0$  from single to double sequences, Acta Mathematica Hungarica 57(1-2) (1991) 129-136.
- [4] F. Moricz and B.E. Rhoades, Almost convergence of double sequences and strong regularity of summability matrices, Mathematical Proceedings of the Cambridge Philosophical Society 104 (1988) 283-294.
- [5] M. Basarir and O. Solancan, On some double sequence spaces, Journal of Indian Academy Mathematics 21(2) (1999) 193-200.
- [6] B.C. Tripathy, On statistically convergent double sequences, Tamkang Journal of Mathematics, 34(3) (2003) 231-237.
- [7] B.C. Tripathy and S. Mahanta, On a class of vector valued sequences associated with multiplier sequences, Acta Mathematica Applicata Sinica (Eng. Ser.) 20(3) (2004) 487-494.
- [8] B.C. Tripathy and M. Sen, Characterization of some matrix classes involving paranormed sequence spaces, Tamkang Journal of Mathematics 37(2) (2006) 155-162.
- [9] B.C. Tripathy and A.J. Dutta, On fuzzy real-valued double sequence spaces  ${}_2\ell_F^p$ , Mathematical and Computer Modelling 46 (9-10) (2007) 1294-1299.

- [10] B.C. Tripathy and B. Hazarika,  $I$ -convergent sequence spaces associated with multiplier sequence spaces, *Mathematical Inequalities and Applications* 11(3) (2008) 543-548.
- [11] B.C. Tripathy and B. Sarma, Statistically convergent difference double sequence spaces, *Acta Mathematica Sinica* 24(5) (2008) 737-742.
- [12] B.C. Tripathy and B. Sarma, Vector valued double sequence spaces defined by Orlicz function, *Mathematica Slovaca* 59(6) (2009) 767-776.
- [13] B.C. Tripathy and H. Dutta, On some new paranormed difference sequence spaces defined by Orlicz functions *Kyungpook Mathematical Journal* 50(1) (2010) 59-69.
- [14] B.C. Tripathy and A.J. Dutta, Bounded variation double sequence space of fuzzy real numbers, *Computers and Mathematics with Applications* 59(2) (2010) 1031-1037.
- [15] B.C. Tripathy and B. Sarma, Double sequence spaces of fuzzy numbers defined by Orlicz function, *Acta Mathematica Scientia* 31 B(1) (2011) 134-140.
- [16] B.C. Tripathy and P. Chandra, On some generalized difference paranormed sequence spaces associated with multiplier sequences defined by modulus function, *Analysis in Theory and Applications* 27(1) (2011) 21-27.
- [17] B.C. Tripathy and A.J. Dutta, Lacunary bounded variation sequence of fuzzy real numbers, *Journal in Intelligent and Fuzzy Systems* 24(1) (2013) 185-189.
- [18] A. Turkmenoglu, Matrix transformation between some classes of double sequences, *Journal of Institute of Mathematics and Computer Science Maths Series* 12(1) (1999) 23-31.
- [19] K. Raj and S.K. Sharma, Lacunary sequence spaces defined by Musielak-Orlicz function, *Le Matematiche* 68 (2013) 33-51.
- [20] K. Raj, A.K. Sharma and S.K. Sharma, Difference sequence spaces in  $n$ -normed spaces defined by a Musielak-Orlicz function, *Armenian Journal of Mathematics* 3 (2010) 127-141.
- [21] K. Raj, A.K. Sharma and S.K. Sharma, A sequence spaces defined by a Musielak-Orlicz function, *International Journal of Pure and Applied Mathematics* 67(4) (2011) 475-484.
- [22] K. Raj and S.K. Sharma, Some sequence spaces in 2-normed spaces defined by Musielak Orlicz function, *Acta Universitatis Sapientiae Mathematica* 3 (1) (2011) 97-109.
- [23] K. Raj A.K. Sharma and S.K. Sharma, Sequence spaces defined by Musielak-Orlicz function in 2-normed spaces, *Journal of Computational Analysis and Applications* 14 (2012).
- [24] K. Raj and S.K. Sharma, Some new Lacunary strong convergent vector-valued multiplier difference sequence spaces defined by a Musielak-Orlicz function *Acta Mathematica Academiae Nyiregyhaziensis* 28 (2012) 103-120.
- [25] K. Raj and S.K. Sharma, Ideal convergent sequence spaces defined by a Musielak-Orlicz function, *Thai Journal of Mathematics* 11 (2013) 577-587.
- [26] J. Lindenstrauss and L. Tzafriri, On Orlicz sequence spaces, *Israel Journal of Mathematics* 10 (1971) 379-390.
- [27] J. Musielak, Orlicz Spaces, *Lectures Notes in Math.*, 1034, Springer-Verlag (1983).
- [28] P.K. Kamthan and M. Gupta, Sequence spaces and series, *Lecture notes in Pure and Applied Mathematics*, 65 Marcel Dekker, Inc., New York (1981).
- [29] V.N. Mishra, Some Problems on Approximations of Functions in Banach Spaces, Ph.D. Thesis, Indian Institute of Technology Roorkee 247 667, Uttarakhand, India (2007).
- [30] V.N. Mishra and L.N. Mishra, Trigonometric Approximation of Signals (Functions) in  $L_p$  ( $p \geq 1$ ) - norm, *International Journal of Contemporary Mathematical Sciences* 7 (19) (2012) 909-918.
- [31] V.N. Mishra, K. Khatri and L.N. Mishra, Strong Cesàro Summability of Triple Fourier Integrals, *Fasciculi Mathematici* 53 (2014).
- [32] L.N. Mishra, V.N. Mishra, K. Khatri and Deepmala, On The Trigonometric approximation of signals belonging to generalized weighted Lipschitz  $W(L^r, \xi(t))$  ( $r \geq 1$ ) - class by matrix  $(C^1.N_p)$  Operator of conjugate series of its Fourier series, *Applied Mathematics and Computation*, Vol. 237 (2014) 252-263.
- [33] V.N. Mishra, K. Khatri, L.N. Mishra, Deepmala; Trigonometric approximation of periodic Signals belonging to generalized weighted Lipschitz  $W'(L_r, \xi(t))$ , ( $r \geq 1$ ) - class by Nörlund-Euler  $(N, p_n)(E, q)$  operator of conjugate series of its Fourier series, *Journal of Classical Analysis*, Volume 5, Number 2 (2014), pp. 91-105. doi:10.7153/jca-05-08.

**Submit your manuscript to IJAAMM and benefit from:**

- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: Articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

---

**Submit your next manuscript at ▶ [editor.ijaamm@gmail.com](mailto:editor.ijaamm@gmail.com)**